

Development of an algorithm for the evaluation of gait and balance impairments in CNS disorders

Ellis RD¹, Kelly PJ¹, Huang C¹
¹ Koneksa Health Inc., New York, NY, USA

Introduction

- Gait and balance is impaired in CNS disorders such as Parkinson's disease (PD)¹ and quantified in surveys e.g. MDS-UPDRS Part III items²
- Gait and balance symptom features can be measured using smartphone applications^{3,4,5,6}
- Koneksa Health developed a new algorithm to calculate gait and balance symptom feature scores (duration, distance, steps, speed, stride period, etc.) and an application for deployment on iPhones
- This study is to determine analytical validity and operational tolerance of this algorithm in healthy subjects using an app deployed on an iPhone 8 plus

Method



- Work under IRB approved study VS001 with 11 self-certified healthy individuals
- Subjects walked a marked course in a large office for each of 5, 10, 15 and 20 seconds under supervision from 3 raters
 - Turns were allowed if a subject walked far enough that they had to turn within space available
- Rater feature scores used as “gold standard” in comparison with corresponding algorithm feature scores
 - Direct measures: duration, steps, distance
 - Derived measures: speed, stride period
- **Analytical validity** was determined by Intraclass Correlation Coefficient (ICC) for agreement between matching algorithm and rater feature scores
 - Classified as excellent ($ICC \geq 0.9$), good ($0.75 \leq ICC < 0.9$), moderate ($0.5 \leq ICC < 0.75$), or poor ($ICC < 0.5$)⁷
- **Operational tolerance** was examined by comparing ICC for the same 20-second walk in
 - Default configuration: placing the phone in a tight trouser pocket
 - Deviations from the default: placing the phone in a loose shorts pocket, and in a shoulder bag

Results

- Total of 83 completed assessments: each subject completed a minimum of 2 attempts at each of the walking test configurations - average subject compliance was 80%
- **Analytical validity:** excellent agreement ($ICC \geq 0.9$) in **duration, distance, and steps** between algorithm and “gold standard” rater feature scores
- **Operational tolerance:** good agreement ($ICC \geq 0.75$) in **distance, steps, stride period** (and speed - except for loose pocket 0.741, close to cutoff)

Reference

1. Grabli, D., et al. (2012). Normal and pathological gait: what we learn from Parkinson's disease. *Journal of Neurology, Neurosurgery & Psychiatry*, 83(10), 979-985.
2. Goetz, C. G., et al. (2008). Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov Disord* 23(15): 2129-2170
3. Bot B. M., et al. (2016), The mPower study, Parkinson disease mobile data collected using ResearchKit. *Scientific Data*. 3:160011
4. Zhan, A., et al., (2018), Using Smartphones and Machine Learning to Quantify Parkinson Disease Severity The Mobile Parkinson Disease Score, *JAMA Neurology* 75(7):876-880
5. Kassavetis, P, et al., (2015), Developing a Tool for Remote Digital Assessment of Parkinson's Disease, *Movement Disorders Clinical Practice* Volume 3, Issue 1:59-64
6. Lipsmeier, F., et al., (2018), Evaluation of smartphone-based testing to generate exploratory outcome measures in a phase 1 Parkinson's disease clinical trial, *Movement Disorders* 33(8): 1287-1297
7. Koo, T. K., & Li, M. Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *Journal of chiropractic medicine*, 15(2), 155-163.



Agreement ICC	Duration	Distance	Steps	Speed	Stride Period
Analytical Validity (combined results for 5s, 10s, 15s, 20s walk)	0.989	0.987	0.992	0.764	0.593
Operational Tolerance (Loose Pocket)	N/A	0.850	0.868	0.741	0.850
Operational Tolerance (Shoulder Bag)	20s walk of fixed duration	0.932	0.858	0.837	0.798

Conclusions

- The new algorithm and iPhone application can accurately measure gait and balance symptom feature scores, i.e. is analytically valid, in healthy individuals under supervision
- The new algorithm is operationally tolerant of different wear modalities
- The new algorithm may be of use in studies of (central nervous system) CNS disorders, such as Parkinson's disease

